

# Genomic prediction is improved

Gert Pedersen Aamand, Lars Peter Sørensen  
og Ulrik Sander Nielsen

November 2016

**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation

STØTTET AF  
mælkeafgiftsfonden

# Observation 1

Larger difference in breeding values than theoretically expected:



(Rel < 90%)




(Rel > 90%)

**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation



# Transition rules – example yield

## Information sources

**Genomic**

**Genomic+daughter**

**Daughter**



**REL < 60%**



**60% < REL < 90%**



**REL > 90%**

**NAV**



**Nordisk Avlsværdi Vurdering** • Nordic Cattle Genetic Evaluation

# Previously: Weight of information

## Information sources

Genomic + ped.

Genomic+daughter

Daughter



REL < 60%



70%



REL > 90%

**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation

# Now: Weight of information (10% polygenic effect)

## Information sources

Genomic + ped.

Genomic+daughter

Daughter



REL < 60%



70%




REL > 90%


**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation




# More smooth transition with polygenic effect





# More smooth transition with polygenic effect



# Introduction af polygenic effect

- All breeds
- All traits
- Both males and females

**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation

# Observation 2

Drop in breeding values from being used for AI to having daughters was larger than expected for young bulls




**Genomic**




**Genomic + first daughter**


# Without standardization




# Standardization – within year




# Standardization – within year



# Standardization – within year + trend across years



# Exempel: NTM for RDC





# Change of NTM

Sum of all changes:

- Improvement of traditional models for fertility, udder, yield and calving
  - Reranking of all animals
- Polygenic effect
  - Especially reranking effect on live genotyped cows and newly proven bulls
- Improved standardization
  - Lower index for genomic tested young AI bulls and genomic tested heifers

**NAV**



# Changes for Holstein bulls born in 2015

Trait	New GEBV	Old GEBV	Difference
<b>Yield</b> (improvements!)	<b>115.1</b>	<b>118.1</b>	<b>-3.0</b>
<b>Milk</b>	<b>108.3</b>	<b>109.9</b>	<b>-1.7</b>
<b>Fat</b>	<b>114.8</b>	<b>116.8</b>	<b>-2.0</b>
<b>Protein</b>	<b>113.8</b>	<b>116.8</b>	<b>-3.1</b>
<b>Growth</b>	<b>99.6</b>	<b>99.6</b>	<b>-0.1</b>
<b>Fertility</b> (improvements!)	<b>108.9</b>	<b>110.6</b>	<b>-1.7</b>
<b>Birth</b> (improvements!)	<b>105.5</b>	<b>106.8</b>	<b>-1.3</b>
<b>Calving</b> (improvements!)	<b>109.2</b>	<b>110.2</b>	<b>-1.1</b>
<b>Udder health</b>	<b>109.4</b>	<b>111.2</b>	<b>-1.8</b>
<b>Other Disease</b>	<b>106.9</b>	<b>107.5</b>	<b>-0.7</b>
<b>Frame</b>	<b>105.7</b>	<b>105.4</b>	<b>0.3</b>
<b>Feet&amp; Legs</b>	<b>108.5</b>	<b>109.1</b>	<b>-0.6</b>
<b>Udder</b> (improvements!)	<b>117.3</b>	<b>117.2</b>	<b>0.1</b>
<b>Milkability</b>	<b>106.7</b>	<b>107.1</b>	<b>-0.5</b>
<b>Temperament</b>	<b>105.2</b>	<b>105.2</b>	<b>0.1</b>
<b>Longevity</b>	<b>114.2</b>	<b>120.8</b>	<b>-6.6</b>
<b>Claw health</b>	<b>107.4</b>	<b>108.3</b>	<b>-0.9</b>
<b>Youngstock survival</b>	<b>101.4</b>	<b>102.8</b>	<b>-1.3</b>
<b>NTM</b>	<b>28.7</b>	<b>33.4</b>	<b>-4.7</b>



# Changes for RDC bulls born in 2015

Trait	New GEBV	Old GEBV	Difference
<b>Yield</b> (improvements!)	<b>112.4</b>	<b>115.3</b>	<b>-2.9</b>
<b>Milk</b>	<b>106.9</b>	<b>109.0</b>	<b>-2.0</b>
<b>Fat</b>	<b>110.8</b>	<b>112.9</b>	<b>-2.1</b>
<b>Protein</b>	<b>111.9</b>	<b>115.0</b>	<b>-3.1</b>
<b>Growth</b>	<b>99.5</b>	<b>99.6</b>	<b>-0.1</b>
<b>Fertility</b> (improvements!)	<b>104.1</b>	<b>104.7</b>	<b>-0.6</b>
<b>Birth</b> (improvements!)	<b>102.6</b>	<b>103.3</b>	<b>-0.7</b>
<b>Calving</b> (improvements!)	<b>103.5</b>	<b>104.8</b>	<b>-1.3</b>
<b>Udder health</b>	<b>107.8</b>	<b>109.1</b>	<b>-1.3</b>
<b>Other Disease</b>	<b>105.4</b>	<b>106.0</b>	<b>-0.6</b>
<b>Frame</b>	<b>103.9</b>	<b>103.9</b>	<b>0.0</b>
<b>Feet&amp; Legs</b>	<b>106.9</b>	<b>108.8</b>	<b>-1.9</b>
<b>Udder</b> (improvements!)	<b>110.3</b>	<b>111.3</b>	<b>-1.0</b>
<b>Milkability</b>	<b>106.9</b>	<b>106.9</b>	<b>0.0</b>
<b>Temperament</b>	<b>102.5</b>	<b>103.3</b>	<b>-0.8</b>
<b>Longevity</b>	<b>111.9</b>	<b>114.4</b>	<b>-3.0</b>
<b>Claw health</b>	<b>102.6</b>	<b>103.0</b>	<b>-0.3</b>
<b>Youngstock survival</b>	<b>99.7</b>	<b>99.6</b>	<b>0.1</b>
<b>NTM</b>	<b>22.1</b>	<b>26.4</b>	<b>-4.3</b>



# Changes for Jersey bulls born in 2015

Trait	New GEBV	Old GEBV	Difference
<b>Yield</b> (improvements!)	<b>111.5</b>	<b>114.1</b>	<b>-2.6</b>
<b>Milk</b>	<b>104.6</b>	<b>105.7</b>	<b>-1.1</b>
<b>Fat</b>	<b>109.5</b>	<b>111.9</b>	<b>-2.4</b>
<b>Protein</b>	<b>110.6</b>	<b>112.3</b>	<b>-1.7</b>
<b>Growth</b>	<b>99.1</b>	<b>99.5</b>	<b>-0.3</b>
<b>Fertility</b> (improvements!)	<b>102.6</b>	<b>102.6</b>	<b>0.1</b>
<b>Birth</b> (improvements!)	<b>100.4</b>	<b>100.0</b>	<b>0.4</b>
<b>Calving</b> (improvements!)	<b>103.3</b>	<b>104.3</b>	<b>-1.0</b>
<b>Udder health</b>	<b>107.2</b>	<b>107.6</b>	<b>-0.4</b>
<b>Other Disease</b>	<b>100.3</b>	<b>100.1</b>	<b>0.2</b>
<b>Frame</b>	<b>106.7</b>	<b>107.6</b>	<b>-0.9</b>
<b>Feet&amp; Legs</b>	<b>104.3</b>	<b>104.3</b>	<b>-0.1</b>
<b>Udder</b> (improvements!)	<b>109.2</b>	<b>108.6</b>	<b>0.5</b>
<b>Milkability</b>	<b>101.7</b>	<b>103.0</b>	<b>-1.3</b>
<b>Temperament</b>	<b>100.1</b>	<b>100.6</b>	<b>-0.5</b>
<b>Longevity</b>	<b>108.1</b>	<b>108.9</b>	<b>-0.8</b>
<b>Claw health</b>	-	-	-
<b>Youngstock survival</b>	-	-	-
<b>NTM</b>	<b>17.3</b>	<b>19.8</b>	<b>-2.5</b>



# Correlation between new and old NTM

	Young Bulls	Heifers	Cows
Holstein	0.95	0.98	0.98
RDC	0.95	0.98	0.98
Jersey	0.96	0.98	0.99

- AI bulls born in 2013-2015
- Cows and heifers born after 2010

**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation

# Changes between new and old NTM

## Genomic tested **bulls** born in year 2015

Change in NTM	Holstein (%)	RDC (%)	Jersey (%)
-10			
-9			
-8	2.2		
-7	3.2	4.1	
-6	18.3	10.2	2.0
-5	33.3	33.7	0.0
-4	26.9	22.4	12.2
-3	8.6	20.4	34.7
-2	7.5	8.2	30.6
-1		1.0	20.4
0			
<b>Average change</b>	<b>-4.7</b>	<b>-4.3</b>	<b>-2.5</b>

**NAV**



Nordisk Avlsværdi Vurdering • Nordic Cattle Genetic Evaluation

# Changes between new and old NTM

## Genomic tested **Cows** born after 2010

Change in NTM	Holstein (%)	RDC (%)	Jersey (%)
-8			
-7	0.2		
-6	1.2	0.1	
-5	4.7	0.6	
-4	12.8	2.9	0.1
-3	22.7	9.8	1.6
-2	26.1	21.2	8.2
-1	20.0	26.9	23.1
0	9.4	22.2	33.2
1	3.3	11.5	24.2
2	0.7	3.8	8.2
3	0.1	0.8	1.3
4		0.1	0.1
5			
<b>Numbers</b>	<b>16,364</b>	<b>20,220</b>	<b>12,704</b>



# Changes between new and old NTM

## Genomic tested **Heifers** born in year 2015

Change in NTM	Holstein (%)	RDC (%)	Jersey (%)
-10			
-9	0.1		
-8	0.9	0.1	
-7	5.5	0.8	
-6	15.3	3.9	0.2
-5	26.8	13.8	1.4
-4	27.6	27.5	10.0
-3	16.5	29.9	29.5
-2	5.9	17.8	34.9
-1	1.3	5.1	19.6
0	0.2	1.0	4.2
1		0.1	0.3
<b>Numbers</b>	<b>10,956</b>	<b>10,218</b>	<b>4,077</b>

**NAV**

